

Regulation of Energy Metabolism by the Extracytoplasmic Function (ECF) σ Factors of *Arcobacter butzleri*

Irati Martinez-Malaxetxebarria^{1,2}, Rudy Muts¹, Linda van Dijk¹, Craig T. Parker³, William G. Miller³, Steven Huynh³, Wim Gaastra¹, Jos P. M. van Putten¹, Aurora Fernandez-Astorga², Marc M. S. M. Wösten^{1*}

1 Department of Infectious Diseases and Immunology, Utrecht University, Utrecht, The Netherlands, **2** Department of Immunology, Microbiology and Parasitology, Faculty of Pharmacy, University of the Basque Country (UPV-EHU), Vitoria-Gasteiz, Spain, **3** Produce Safety and Microbiology Research Unit, Agricultural Research Service, US Department of Agriculture, Albany, California, United States of America

Abstract

The extracytoplasmic function (ECF) σ factors are fundamental for bacterial adaptation to distinct environments and for survival under different stress conditions. The emerging pathogen *Arcobacter butzleri* possesses seven putative pairs of σ /anti- σ factors belonging to the ECF family. Here, we report the identification of the genes regulated by five out of the seven *A. butzleri* ECF σ factors. Three of the ECF σ factors play an apparent role in transport, energy generation and the maintenance of redox balance. Several genes like the *nap*, *sox* and *tct* genes are regulated by more than one ECF σ factor, indicating that the *A. butzleri* ECF σ factors form a network of overlapping regulons. In contrast to other eubacteria, these *A. butzleri* ECF regulons appear to primarily regulate responses to changing environments in order to meet metabolic needs instead of an obvious role in stress adaptation.

Citation: Martinez-Malaxetxebarria I, Muts R, van Dijk L, Parker CT, Miller WG, et al. (2012) Regulation of Energy Metabolism by the Extracytoplasmic Function (ECF) σ Factors of *Arcobacter butzleri*. PLoS ONE 7(9): e44796. doi:10.1371/journal.pone.0044796

Editor: Roy Martin Roop II, East Carolina University School of Medicine, United States of America

Received: January 19, 2012; **Accepted:** August 14, 2012; **Published:** September 18, 2012

This is an open-access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the Creative Commons CC0 public domain dedication.

Funding: This work was supported in part by the Basque Government (IT-528-10) through a Predoctoral Fellowship of the Basque Government for the Training of Researches (BF106.85); by a parallel incentive within the Predoctoral Fellowship of the Basque Government for short stays in different Research Centers (EC-2009-1-4); and by a European Molecular Biology Organization (EMBO) Short Term Fellowship (ASTF29-2010) to IMM. This work was also supported by a fellowship of Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO-Vidi) grant 917.66.330 to MMSMW and by funds from US Department of Agriculture Current Research Information System (USDA-CRIS) project 5325-42000-047-00D. No additional external funding received for this study. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: M.Wosten@uu.nl

Introduction

Arcobacter spp. are Gram-negative, small, motile and spiral-shaped bacteria belonging to the *Campylobacteraceae* family. *Arcobacter* is currently comprised of fifteen species [1,2]. Within this genus, *A. butzleri*, *A. cryaerophilus* and *A. skirrowii* are associated with animal and human diseases, such as reproductive disorders, mastitis and diarrhoea in animals [3–6], and enteritis and occasionally bacteraemia in humans [7–10]. *Arcobacter* spp. have been classified in 2002 as emerging pathogens by the ICMSF (International Commission on Microbiological Specifications for Foods) [11]. The main reservoirs for *Arcobacter* spp. are water environments, especially sewage and coastal waters [12–14].

In 2007 the complete genome of the human clinical isolate *A. butzleri* RM4018 was sequenced [15]. The bacterium appears to have a large number of signal transduction systems, indicating that this organism is able to respond to many different environmental signals. Apart from having one of the highest densities of two component systems per megabase, *A. butzleri* RM4018, unlike *Campylobacter* spp. or *Helicobacter* spp., is also predicted to contain seven σ /anti- σ factor pairs belonging to the ECF family (Extracytoplasmic function) of sigma factors. The seven *A. butzleri* RM4018 ECF σ factor pairs are genetically unlinked but for each pair, the σ factor gene is positioned adjacent to and in the same

orientation as the putative cognate anti- σ factor gene. The amino acid identity between any two of the seven ECF σ factors varies between 30 and 49%, suggesting that each ECF pair plays a different role in the biology of strain RM4018 and responds perhaps to a different suite of external signals.

ECF σ factors have been identified in both Gram-negative and Gram-positive bacteria and they are fundamental for bacterial adaptation to different environments and for survival under different stress conditions [16,17]. Their role in cellular physiology is highly variable and includes adaptation to membrane-affecting compounds, extreme temperatures and pH, light, high pressure, carbon, nitrogen and iron starvation, oxidative and osmotic stress, and the regulation of virulence in pathogenic organisms [16,18]. Despite their diverse effects, ECF σ factors share common features. In most cases ECF σ factors are co-transcribed with their cognate anti- σ factor. The anti- σ factor, which often spans the inner membrane, sequesters and inactivates the cognate σ factor at the inner face of the cytoplasmic membrane. Upon a specific external stimulus, the ECF anti- σ factor is inactivated by protein degradation and the σ factor is released from the membrane, where it associates with RNA polymerase and mediates transcriptional activation of its target genes [19]. The ECF anti- σ factor can also be released from the ECF σ factor by a phosphorylated response regulator NepR, this partner-switching mechanism has

recently be discovered in alphaproteobacteria [20]. Some ECF σ /anti- σ factors form together with an outer membrane TonB-dependent receptor, a trans-envelope signal transduction pathway [31]. These TonB-dependent receptors are involved in the uptake of specific molecules but also they sense and transmit, via the anti- σ factor, extracellular signals, which lead to the activation of a specific ECF σ factor.

In the present study, we investigated the role of the *A. butzleri* RM4018 ECF σ factors. We developed the genetic tools to manipulate *A. butzleri* and determined the regulons of most ECF σ factors using a combination of microarray-based transcriptome analysis and functional assays.

Materials and Methods

Bacterial Strains, Plasmids and Growth Conditions

Bacterial strains and plasmids used in this work are listed in Table 1. *A. butzleri* strains were routinely grown at 30°C in Brain Heart Infusion (BHI) broth (Oxoid) or on Mueller Hinton (MH) agar (Oxoid) supplemented with 5% sheep blood (Biotrading). *E. coli* strains were routinely grown at 37°C in Luria-Bertani (LB) broth or on LB agar plates (Biotrading) supplemented with ampicillin (100 μ g/ml) or kanamycin (50 μ g/ml) when needed.

Construction of A. Butzleri ECF σ Factor and ECF Anti- σ Factor Mutants

The genes encoding the seven ECF σ factors and their cognate anti- σ factors were amplified using the primer pairs AB0986F/AB0987R, AB1044F/AB1044R, AB1437F2/AB1437R2, AB1460F/AB1460R, AB1576F/1576R, AB2151F/AB2151R and AB2300F/2300R (Table 2), the proofreading enzyme *Pfu* (Promega) and *A. butzleri* RM4018 chromosomal DNA as template. The PCR products were tailed with a 5'-A nucleotide using *Taq* polymerase (Invitrogen) and ligated into the pGEM-T Easy vector (Promega) to obtain pGEMab0983–0984, pGEMab1040–1041, pGEMab1429–1430, pGEMab1452–1453, pGEMab1567–1568, pGEMab2164–2165 and pGEMab2315–2316 (Table 1). Inverse PCR was performed on the σ /anti- σ plasmids to delete a large part of both the σ and anti- σ factor (σ /anti- σ knock-out plasmids) or of only the anti- σ factor encoding genes (anti- σ knock-out plasmids). Unique *Bam*HI restriction sites were introduced at the same time. The plasmid pGEM Δ ab1429–1430 was obtained by digesting plasmid pGEM Δ ab1429 with *Bam*HI and *Bcl*I. The knockout constructs were created by digestion of the inverse PCR products with *Bam*HI and ligation to a 1.4-kb *Bam*HI fragment containing a kanamycin resistance gene (*aph*(3')-III) from pMW2 [21].

The ECF σ factor genes were inactivated by marker exchange mutagenesis. First, the knock-out plasmids were introduced into *A. butzleri* RM4018 by electrotransformation. To obtain electrocompetent *A. butzleri* RM4018, a 5 ml overnight culture was diluted 20 times in 100 ml fresh BHI medium and incubated at 30°C on a shaking platform (150 rpm). Bacteria were harvested by centrifugation (4,500 \times g, 1 h, 4°C) when the optical density at 550 nm had reached values between 0.2 and 0.6. The bacteria were washed twice in 5 ml of ice-cold sucrose-glycerol solution (15% glycerol; 272 mM sucrose in water), resuspended in 0.5 ml ice-cold sucrose-glycerol solution and aliquoted into 50 μ l solutions containing approximately $3-5 \times 10^9$ CFU. One μ g of each plasmid was added to 50 μ l (approximately $3-5 \times 10^9$ CFU) of competent *A. butzleri* RM4018 and incubated for 3 min on ice. The cells were transferred to a 0.2 cm electroporation cuvette (Bio-Rad) and electroporated using a Bio-Rad Gene Pulser set at 2.25 kV, 400 Ω and 25 μ F. Bacteria were recovered in 1 ml of BHI broth for

10 min at room temperature, transferred to 2 ml of pre-warmed BHI and incubated for 3 hours at 30°C on a shaking platform (150 rpm). Mutants were selected by growing the cells for two to five days at 30°C aerobically on kanamycin-containing MH plates. Homologous recombination resulting in double-crossover events was verified by PCR.

Phenotype Characterization

Growth curves were generated by diluting pre-cultures grown overnight in BHI to a starting optical density (OD₅₅₀) of 0.05 in 30 ml of BHI. The 30 ml cultures were grown in conical flask under aerobic conditions, 150 rpm at 30°C. Bacterial growth and cell density were monitored by measuring the absorbance at 550 nm at different time intervals. The exponential growth rate was calculated from four separated growth experiments. Several stress conditions which limited the growth of the wildtype *Arcobacter* strain were tested. To measure the influence of extreme temperatures, the starting cultures were incubated at 4°C or 60°C for 15 min prior to their incubation at 30°C. To study the effect of the pH on the growth of the strains, the pH of the BHI was adjusted to pH 5 or 9. Osmotic stress was tested by adding 0.35 M NaCl to the BHI. In additional experiments, other stress-inducing chemicals were added to the BHI to concentrations which limited the growth of the wildtype *Arcobacter* strain. These chemicals included: ethanol (5%); SDS (1%); the iron chelator 2,2-dipyridyl (300 μ M); the oxidative stress-inducing chemicals H₂O₂ (0.04%) and diamide (2 mM); or antimicrobial compounds such as penicillin G and polymyxin B (25 μ g/ml). Motility assays were performed by stabbing the strains with a pipette tip into semisolid medium (thioglycolate medium containing 0.4% agar, Difco), followed by incubations under aerobic conditions at 30°C or 37°C for 48 hours.

RNA Isolation

Overnight grown cultures of *A. butzleri* were diluted to an OD₅₅₀ of 0.1 in BHI and incubated at 30°C on a shaking platform set at 150 rpm. RNA was isolated from 5 ml of mid-logarithmic phase cultures (OD₅₅₀ of approximately 0.5), using the RNA-Bee™ kit (Tel-Test, Inc) following the manufacturers specifications.

Microarray Hybridization and Analysis

For expression profiling, an indirect comparison of gene expression levels was performed [22,23]. In this microarray experimental design, each labeled cDNA was combined with labeled genomic DNA from *A. butzleri* RM4018. Mixtures were hybridized to a previously designed and manufactured *A. butzleri* DNA array [15]. Labeling of RNA and DNA, hybridization procedure and microarray data analysis were performed as previously described [24]. Details of the microarray have been deposited in the NCBI GEO repository (<http://www.ncbi.nlm.nih.gov/geo/>) under platform accession number GPL14948. The microarray data set has been deposited in the NCBI Gene Expression Omnibus (GEO) repository (<http://www.ncbi.nlm.nih.gov/geo/>) under accession number GSE34089.

Real-time RT-PCR

Primers used in this assay were designed using the Primer Express software (Applied Biosystems) and are listed in Table 2. Prior to amplification, RNA samples were treated with RNase-free DNase I (Invitrogen). RT-PCR was performed on 0.2 μ g of DNase I treated RNA with 1 μ M of primers and the SYBR® Green I kit (Eurogentec) using a LightCycler® 480 Real-Time PCR System (Roche). The PCR parameters were 30 min at 48°C;

Table 1. Bacterial strains and plasmids used in this study.

Bacterial strain or plasmid	Origin/function	Source
A. butzleri strains		
<i>A. butzleri</i> RM4018	Human clinical isolate (ATCC 49616)	USDA*
<i>A. butzleri</i> $\Delta\sigma^1/A\sigma^1::Km$	RM4018 derivative $\Delta ab0983-0984::aph(3')-III$	This study
<i>A. butzleri</i> $\Delta A\sigma^1::Km$	RM4018 derivative $\Delta ab0984::aph(3')-III$	This study
<i>A. butzleri</i> $\Delta\sigma^2/A\sigma^2::Km$	RM4018 derivative $\Delta ab1040-1041::aph(3')-III$	This study
<i>A. butzleri</i> $\Delta A\sigma^2::Km$	RM4018 derivative $\Delta ab1041::aph(3')-III$	This study
<i>A. butzleri</i> $\Delta\sigma^4/A\sigma^4::Km$	RM4018 derivative $\Delta ab1452-1453::aph(3')-III$	This study
<i>A. butzleri</i> $\Delta A\sigma^4::Km$	RM4018 derivative $\Delta ab1453::aph(3')-III$	This study
<i>A. butzleri</i> $\Delta\sigma^5/A\sigma^5::Km$	RM4018 derivative $\Delta ab1567-1568::aph(3')-III$	This study
<i>A. butzleri</i> $\Delta A\sigma^5::Km$	RM4018 derivative $\Delta ab1567::aph(3')-III$	This study
<i>A. butzleri</i> $\Delta\sigma^7/A\sigma^7::Km$	RM4018 derivative $\Delta ab2315-2316::aph(3')-III$	This study
<i>A. butzleri</i> $\Delta\sigma^7::Km$	RM4018 derivative $\Delta ab2315::aph(3')-III$	This study
E. coli strains		
<i>E. coli</i> DH5 α NCCB2955	Competent cells for cloning	NCCB**
Plasmids		
pGEM-T Easy	Cloning vector, Amp ^r	Promega
pGEMab0983-0984	pGEM-T Easy containing <i>ab0983-0984</i>	This study
pGEMab1040-1041	pGEM-T Easy containing <i>ab1040-1041</i>	This study
pGEMab1429-1430	pGEM-T Easy containing <i>ab1429-1430</i>	This study
pGEMab1452-1453	pGEM-T Easy containing <i>ab1452-1453</i>	This study
pGEMab1567-1568	pGEM-T Easy containing <i>ab1567-1568</i>	This study
pGEMab2164-2165	pGEM-T Easy containing <i>ab2164-2165</i>	This study
pGEMab2315-2316	pGEM-T Easy containing <i>ab2315-2316</i>	This study
pGEM $\Delta ab0983-0984$	pGEM-T Easy containing $\Delta ab0983-0984$	This study
pGEM $\Delta ab1040-1041$	pGEM-T Easy containing $\Delta ab1040-1041$	This study
pGEM $\Delta ab1429-1430$	pGEM-T Easy containing $\Delta ab1429-1430$	This study
pGEM $\Delta ab1452-1453$	pGEM-T Easy containing $\Delta ab1452-1453$	This study
pGEM $\Delta ab1567-1568$	pGEM-T Easy containing $\Delta ab1567-1568$	This study
pGEM $\Delta ab2164-2165$	pGEM-T Easy containing $\Delta ab2164-2165$	This study
pGEM $\Delta ab2315-2316$	pGEM-T Easy containing $\Delta ab2315-2316$	This study
pGEM $\Delta ab0984$	pGEM-T Easy containing $\Delta ab0984$	This study
pGEM $\Delta ab1041$	pGEM-T Easy containing $\Delta ab1041$	This study
pGEM $\Delta ab1429$	pGEM-T Easy containing $\Delta ab1429$	This study
pGEM $\Delta ab1453$	pGEM-T Easy containing $\Delta ab1453$	This study
pGEM $\Delta ab1568$	pGEM-T Easy containing $\Delta ab1568$	This study
pGEM $\Delta ab2164$	pGEM-T Easy containing $\Delta ab2164$	This study
pGEM $\Delta ab2316$	pGEM-T Easy containing $\Delta ab2316$	This study
pGEM $\Delta ab1040-1041::km$	pGEM-T Easy containing $\Delta ab1040-1041::aph(3')-III$	This study
pGEM $\Delta ab1429-1430::km$	pGEM-T Easy containing $\Delta ab1429-1430::aph(3')-III$	This study
pGEM $\Delta ab1452-1453::km$	pGEM-T Easy containing $\Delta ab1452-1453::aph(3')-III$	This study
pGEM $\Delta ab1567-1568::km$	pGEM-T Easy containing $\Delta ab1567-1568::aph(3')-III$	This study
pGEM $\Delta ab2164-2165::km$	pGEM-T Easy containing $\Delta ab2164-2165::aph(3')-III$	This study
pGEM $\Delta ab2315-2316::km$	pGEM-T Easy containing $\Delta ab2315-2316::aph(3')-III$	This study
pGEM $\Delta ab0984::km$	pGEM-T Easy containing $\Delta ab0984::aph(3')-III$	This study
pGEM $\Delta ab1041::km$	pGEM-T Easy containing $\Delta ab1041::aph(3')-III$	This study
pGEM $\Delta ab1429::km$	pGEM-T Easy containing $\Delta ab1429::aph(3')-III$	This study
pGEM $\Delta ab1453::km$	pGEM-T Easy containing $\Delta ab1453::aph(3')-III$	This study
pGEM $\Delta ab1568::km$	pGEM-T Easy containing $\Delta ab1568::aph(3')-III$	This study
pGEM $\Delta ab2164::km$	pGEM-T Easy containing $\Delta ab2164::aph(3')-III$	This study

Table 1. Cont.

Bacterial strain or plasmid	Origin/function	Source
pGEM Δ ab2316::km	pGEM-T Easy containing Δ ab2316::aph(3')-III	This study
pMW2	Plasmid with kanamycin resistance (aph(3')-III) gene from <i>Campylobacter</i> .	[21]

*USDA, Agricultural Research Service, California, USA.

**The Netherlands Culture Collection of Bacteria.

doi:10.1371/journal.pone.0044796.t001

10 min at 95°C, 40 cycles of 95°C for 15 s and 60°C for 1 min and a final dissociation step of 95°C for 15 s, 60°C for 20 s and 95°C for 15 s. To compensate for the variance in the amount of mRNA in the reactions, the calculated threshold cycle (C_t) for each gene amplification was normalized to the C_t values for the *gyrA* gene. Fold changes were calculated according to the $\Delta\Delta C_t$ method [25]. Each sample was examined in four replicates and was repeated with at least two independent preparations of RNA. Standard deviations were calculated and displayed as error bars.

Nitrate/nitrite Assay

The nitrate reductase activity was determined by measuring the production of nitrite from nitrate as previously described [26]. Briefly, strains were grown overnight (16 h) under aerobic conditions in BHI broth containing 20 mM of sodium nitrate. Nitrite accumulation in the supernatant was detected by mixing 50 μ l of the culture supernatants with 850 μ l of 1% (w/v) sulphanilamide dissolved in 1 M of HCl and 100 μ l of 0.02% (w/v) naphthylethylenediamine solution. After 15 minutes the formation of p-sulfobenzene-azo α -naphthylamine was measured at 540 nm. The amount of nitrite present in 50 μ l culture supernatant was estimated using a nitrite standard curve. The nitrite production was adjusted to the total bacterial proteins present in the culture as estimated by using the BCA protein assay kit. To determine the nitrate concentration in *Arcobacter* culture supernatants the supernatants were 10 times diluted in BHI. The diluted culture supernatants were incubated with sulphanilamide and naphthylethylenediamine as described for the nitrite detection. Next a trace amount of zinc dust was added to the mixture. The zinc dust reduces nitrate to nitrite and a red color develops due to the formation of p-sulfobenzene-azo α -naphthylamine. After 2 minutes the zinc has settled on the bottom and 900 μ l of the supernatant was transferred to a new tube and the absorbance of the red color was measured at 540 nm. The amount of nitrate present in 50 μ l culture supernatant was estimated using a nitrate standard curve which was also treated with zinc dust. Reactions were measured four times and repeated with two independent cultures.

Sox Enzyme Activity Assay

Overnight cultures in BHI were washed once with *Arcobacter* Minimal 1 (AM1) medium (4.2 mM Na₂HPO₄, 2.4 mM KH₂PO₄, 9 mM NaCl, 19 mM NH₄Cl 1 \times M9 salts, 1 mM of MgCl₂, 0.2% of sodium pyruvate, 0.02% of FeNH₄citrate; adjusted to pH 7.5) amended with 20 mM of sodium sulfite, and 10 μ g of phenol red per milliliter. The bacteria were subsequently diluted in sodium sulfite-, phenol red-amended AM1 medium to an OD₅₅₀ of 0.05 and allowed to grow in completely filled capped 5 cm glass vials. The strains which were able to oxidize sulfur caused a drop in the pH that was visible as a color change from red to yellow. After 6 hour of growth at 30°C, absorbance at 564 nm was measured and the results obtained for the wild-type strain were set at 100%.

Citrate Uptake Assay

The ability of the strains to utilize citrate was determined using Simmons citrate agar (Oxoid). Strains grown overnight on MH agar supplemented with 5% sheep blood were streaked onto Simmons Citrate Agar supplemented with 0.01% sodium pyruvate. Cells were incubated at 30°C under aerobic conditions for 48 hours. The strains which were able to use the citrate as carbon source increased the pH of medium, resulting in a color change of the medium from green to blue.

Statistical Analyses

Analysis of variance (ANOVA) was used to identify if any of the mutations significantly changed the growth rate of the *Arcobacter* strains. Statistical analyses were performed using the SPSS 19.0 statistical package program (SPSS Inc., Chicago, IL).

Results

Mutagenesis of the ECF σ Factor and Anti- σ Factor Encoding Genes

For each ECF σ /anti- σ pair, mutational inactivation of the anti- σ factor results in constitutive induction of the σ response, while mutational inactivation of the σ factor results in the down-regulation of ECF-dependent transcription [19,27]. This allowed us to study the function of the ECFs in *A. butzleri* strain RM4018 without knowing the specific signal(s) that activate them. In order to address the role of the seven ECF σ factors in *A. butzleri* RM4018, we inactivated either both the σ and the anti- σ factor, or the anti- σ factor alone. Since no suitable *Arcobacter* antibiotic resistance cassettes were available and the previously-used chloramphenicol cassette of *Campylobacter coli* could not be used since *A. butzleri* RM4018 is resistant to chloramphenicol, we replaced a large part of the ECF coding regions with a kanamycin resistance gene (*aph(3')-III*) of *C. coli* [21]. After modifying the *Arcobacter* mutagenesis protocol described by Ho *et al.* [28], we were able to replace the ECF σ /anti- σ factor genes *AB0986/0987* (σ^1 , A σ^1), *AB1044/1045* (σ^2 , A σ^2), *AB1460/1461* (σ^4 , A σ^4), *AB1576/1577* (A σ^5 , σ^5) and *AB2300/2301* (A σ^7 , σ^7) with the Km^r cassette. We also obtained single mutants in the anti- σ factor genes A σ^1 , A σ^2 , A σ^4 , A σ^5 and A σ^7 . All mutants contained the Km^r cassette in the same orientation as the ECF genes. As no suitable *Arcobacter* shuttle plasmids exist and no other antibiotic cassettes could be used, we were unable to perform complementation studies.

Phenotype Characterization

Comparison of the various mutants with the parent strain RM4018 revealed neither differences in bacterial shape or colony formation on sheep blood plates nor in growth rate in BHI medium as statistically estimated by Anova F = 1,777 (P > 0.05) (Figure 1A and B). A common role of ECF σ factors is to protect bacteria against external stress [16]. Exposure of the *A. butzleri* mutant strains to extreme temperatures (e.g., 4°C or 60°C), pH 5

Table 2. Oligonucleotides used in this study.

Primer name	DNA sequence (5' → 3')
Mutant construction	
AB0986F	AACAGTTGCTGATATTAAGCTATG
AB0986R	TCTTTTATACCATCAAAATGCTGC
AB0986F-bamHI	<u>AGGATCCT</u> GATATTTGAAGATGAAAAATCTCTCA
AB0986R-bamHI	<u>AGGATCCAGT</u> TTCATTTTCTTCAAAAGG
Arcoant σ 1-bamHI	<u>AGGATCCGTT</u> CAAAAGAGGTTAAACCTC
AB1044F	TAAGTTTAGTTTCAAAATCTCTGA
AB1044R	AACACTATCTCCAAATTTGATATC
AB1044F-bamHI	<u>TGGATCCG</u> AAAAAGTATCTAAAACGGAA
AB1044R-bamHI	<u>TGGATCCT</u> ATTGGTGAGATTTTATCTATGT
Arcoant σ 2-bamHI	<u>AGGATCCCT</u> AACCACTCTTTAATTCAATT
AB1437F2	CTTATGATTTTTATAAATATATATGTA
AB1437R2	ATGCTTAAATATTATGATGAGTT
AB1437R-bamHI	<u>AGGATCCCT</u> ATTCTAACATCTGGTAAATG
Arcoant σ 3-bamHI	<u>AGGATCCT</u> CATGTAATAAAATAAGCTGC
AB1460F	ACTATTCCAATTATTACAGCATT
AB1460R	AGAAAACTCTTATTACATTGTAA
AB1460F-bamHI	<u>AGGATCCAGT</u> GGAATTTTGAAGTC
AB1460R-bamHI	<u>CGGATCCT</u> TGTGCATTATCTTTGTTGG
Arcoant σ 4-bamHI	<u>AGGATCCT</u> GGTAAGGCTTTTATTGAGTT
AB1576F	TGCTGTAAAGTTTACAATAAAGAA
AB1576R	TATTCCATCTTCAAGAGCTT
AB1576F-bamHI	<u>AGGATCCT</u> GCTCTTTGGTTTCAATT
AB1576R-bamHI	<u>TGGATCCG</u> TGAATTTTCATCACTCA
Arcoant σ 5-bamHI	<u>AGGATCCAT</u> ATGTTGCTTCTTCTTTGATT
AB2151F	TTCTAATACTGTTCTTGACC
AB2151R	TTTTACAAGAGCATTATATCG
AB2151F-bamHI	<u>TGGATCCT</u> ACAAGTTTGGAAACATAG
AB2151R-bamHI	<u>TGGATCCG</u> AAAAACAGATATAGAAGAGA
Arcoant σ 6-bamHI	<u>TGGATCCT</u> CCACATAAAATAGCAGCCATT
AB2300F	ACAATCAAGTTTTCGCGCA
AB2300R	CTTAATAAAGCCACTCCTTG
AB2300F-bamHI	<u>AGGATCCT</u> CATGGGCTAAATCTTTGTC
AB2300R-bamHI	<u>AGGATCCG</u> TGAACAAATACCAGAAAATA
Arcoant σ 7-bamHI	<u>AGGATCCA</u> CAACTCAACCAATATCTTGCTT
qRT-PCR	
arconapBtaqF	TGACCACTTATCAAATGCGAGATT
arconapBtaqR	GGAACATTTGCACTATTTGATTGTG
arconapAtaqF	GCCGCTGCGGCTGTT
arconapAtaqR	CGCAACCGCTTCTG
arcsoxDtaqF	TGCGCAATGTGTTATGTGTCA
arcsoxDtaqR	GAACCACCGACAAGCATAGGA
arcsoxAtaqF	CAGTGCATAGACTAAGATGGGAAGAA
arcsoxAtaqR	GCAACTTGCCTTGGTCAACAA
Abu100tctAtaqF	GGATTAGTGCAACCCGAGAGT
Abu100tctAtaqR	CGTCCACTTCTGGAATTCC
Abu98tctCtaqF	ACAGCAGCAAGACAACAAAATACTTT
Abu98tctCtaqR	AAGATTGAGGAAATACACCTTGTAAATGA
AB0988Ftaq	ATAAAAGAGGTAGTGTGAAAGTGTT

Table 2. Cont.

Primer name	DNA sequence (5' → 3')
AB0988Rtaq	CGGTTGATGTGGCTGTAAATTC
AB1462Ftaq	TTCCAAGAAATCCAATGAACATG
AB1462Rtaq	AAGCACTCCATTGGTCATTAATACTATAT
AB1573Ftaq	ACCATAACTTATACTAGCATTCCCTTTTG
AB1473Rtaq	GTACGTGGAGCAACAGGACTAACTAC
arcogyrAtaqF2	AAAGTTTCTCTTAAAGAGCTCCACTTG
arcogyrAtaqR2	TGTATATCTCATAGCCGCTGCATT

*Bam*HI restriction sites are underlined.
doi:10.1371/journal.pone.0044796.t002

or 9, or sub-inhibitory concentrations of stress-inducing chemicals (e.g., ethanol, SDS, metal ion chelators EDTA or 2,2-dipyridyl, the oxidative stress-inducing chemicals H₂O₂ and diamide, or the antimicrobial compounds penicillin G and polymyxin B) did not yield significant growth or morphology differences (data not shown). Similarly, the motility of the mutants on semisolid medium was unchanged compared to the wild-type. These results indicated that the functions of the *Arcobacter* ECF σ factors might be different from those in other bacterial species.

Identification of ECF σ Factor-regulated Genes

To identify the genes regulated by the RM4018 ECF σ factors, RNA was isolated from wild-type and mutant strains grown to mid logarithmic phase and subjected to microarray-based transcriptome analysis. To obtain maximal ECF σ factor-dependent transcript differences, transcripts of each $\sigma/\Delta\sigma$ mutant and its cognate $\Delta\sigma$ mutant were compared. Genes showing more than a fourfold change in transcript levels were considered as ECF σ -dependent. Based on this criterion, the ECF σ factors σ^1 , σ^2 , σ^4 , σ^5 and σ^7 were found to regulate 3, 65, 14, 42 and 72 genes, respectively (Tables S1 to S5). None of the ECF σ factors appeared to regulate its own transcription. Interestingly, more than thirty genes were regulated by two or more ECF σ factors (Table 3). In agreement with the phenotypic characterization, the identified ECF σ -dependent genes indicate that the ECF σ factors of *A. butzleri* regulate other genes than ECF σ factors in other bacterial species.

Genes Regulated by AB0986 (ECF σ^1)

The transcriptome analysis revealed that σ^1 activates the transcription of *AB0988*, encoding a putative TonB-dependent receptor protein, and down-regulates the transcription of *AB1593* and *AB0033*, that encode a putative sodium: alanine symporter and a L-lactate permease, respectively (Table S1).

Genes Regulated by AB1044 (ECF σ^2)

Based on the microarray data, ECF σ^2 activates the transcription of 35 genes and caused a down-regulation of the expression of 30 genes (Table S2). All annotated gene products represent putative proteins. The sodium: solute symporter AB0504 is the most strongly up-regulated gene product. Apart from this, ECF σ^2 is predicted to control the transcription of the genetic loci: *AB0345-AB0346*, encoding a Nrf-type nitrite reductase; *AB0353* to *AB0359*, coding for a Nap- type nitrate reductase and a C₄-dicarboxylate transport system; *AB0376-0377*, encoding an aldehyde dehydrogenase and an uncharacterized protein; *AB0494-AB0495*, encoding an acetate kinase and a phosphate




Figure 1. Arcobacter growth curves and rates. (A) Growth curves of the wild-type strain RM4018, the ECF anti- σ mutants and the ECF σ /anti- σ double mutants. Strains were grown in BHI at 30°C under aerobic conditions. Data correspond to one representative of four independent experiments. (B) Exponential growth rates (hr⁻¹) with standard deviations of four growth curves including one shown in (A). doi:10.1371/journal.pone.0044796.g001

acetyltransferase; *AB0563* to *AB0570*, encoding the Sox multi-enzyme complex; and *AB1442–1443*, encoding a Ni/Fe-hydrogenase. A putative operon (*AB0102–0106*) coding for the tricarboxylic transport proteins TctABC and a large locus (*AB1664* to *AB1682*) coding for phage related proteins were among the genes that are down-regulated by ECF σ^2 .

Genes Regulated by AB1460 (ECF σ^4)

According to the transcriptomics data, σ^4 activates 11 genes and caused a down-regulation of the expression of 3 genes (Table S3). The main up-regulated gene product is the putative TonB-dependent receptor protein (AB1462) encoded by the same locus encoding the σ^4 /anti- σ^4 factors. Among the other genes up-regulated by ECF σ^4 are: *napHGA* (*AB0354* to *AB0356*), *aceEF* (*AB1480–1481*), *frdA* (*AB0297*), *nrfA* (*AB0345*) and *filhA1* (*AB1507*) encoding, respectively, a part of a Nap-type nitrate reductase, two pyruvate dehydrogenase components, a fumarate reductase subunit, a cytochrome *c*₅₅₂ nitrate reductase and a formate dehydrogenase subunit. A methyl-accepting chemotaxis protein (AB0602) is the most strongly down-regulated gene product.

Genes Regulated by AB1577 (ECF σ^5)

Identification of ECF σ^5 -dependent genes by transcriptomics analysis revealed 16 up-regulated and 26 down-regulated genes (Table S4). The closely-linked TonB-dependent receptor gene *AB1573*, was the most strongly up-regulated. The putative TctABC transport proteins AB0102–AB0104, the carbonic anhydrase AB0107, the putative ammonia monooxygenase AB0108, and the ubiquinolcytochrome *c* oxidoreductase encoded by the *petAB* (*AB2054–AB2055*) genes are among the gene products up-regulated by σ^5 . The genes *fur2* and *cspA*, and the genetic locus *AB1062–1066*, which includes NADPH quinoreductase, NADPH:flavodoxin oxidoreductase, and NADPH nitroreductase, were found to be repressed.

Genes Regulated by AB2301 (ECF σ^7)

The transcriptome analysis revealed that σ^7 activates 37 genes and represses another 35 (Table S5). Almost all annotated genes repressed by σ^7 were activated by σ^2 . The genes encoding the putative TctABC transport proteins (*AB0103–0105*) are the most up-regulated genes by σ^7 , while the *sox* operon and *AB0494–AB0495*, encoding an acetate kinase and a phosphate acetyltrans-

Table 3. Genes regulated by two or more ECF σ factors.

Functional class ^a	ORF ^a	Gene ^a	Transcription activated by	Transcription repressed by
Energymetabolism: anaerobic respiration	AB0345	<i>nrfA</i>	σ^2 and σ^4	
	AB0354	<i>napH</i>	σ^2 and σ^4	
	AB0355	<i>napG</i>	σ^2 and σ^4	
	AB0356	<i>napA</i>	σ^2 and σ^4	
	AB0297	<i>frdA</i>	σ^4 and σ^5	
Energymetabolism: aerobic respiration.	AB1442	<i>hyaB</i>	σ^2 and σ^4	σ^7
Energymetabolism: pyruvate dehydrogenase.	AB1481	<i>aceF</i>	σ^4 and σ^5	
Energymetabolism: electrontransport.	AB2055	<i>petA</i>	σ^5	
Transport/binding proteins: carbohydrates, organic acids and alcohols	AB0357	<i>dctP</i>	σ^2 and σ^7	σ^7
	AB0358	<i>dctQ</i>	σ^2	σ^7
	AB0359	<i>dctM</i>	σ^2	σ^7
	AB0102		σ^5 and σ^7	σ^2
	AB0103		σ^5 and σ^7	σ^2
	AB0104		σ^5 and σ^7	σ^2
Transport/binding proteins: others	AB0504		σ^2 and σ^5	σ^7
Central intermediary metabolism: general	AB0376	<i>ald</i>	σ^2	σ^7
	AB0107	<i>cynT1</i>	σ^5	σ^7
Central intermediary metabolism: sulfur metabolism.	AB0563	<i>soxC</i>	σ^2	σ^7
	AB0564	<i>soxD</i>	σ^2	σ^7
	AB0565	<i>soxX</i>	σ^2	σ^7
	AB0566	<i>soxY</i>	σ^2	σ^7
	AB0567	<i>soxZ</i>	σ^2	σ^7
	AB0568	<i>soxA</i>	σ^2	σ^7
	AB0569		σ^2	σ^7
	AB0570	<i>soxB</i>	σ^2	σ^7
Small molecule metabolism: carbon compound degradation.	AB0494	<i>ackA1</i>	σ^2	σ^7
	AB0495	<i>pta</i>	σ^2	σ^7
Cell processes: detoxification.	AB1553	<i>katG</i>	σ^2	σ^7
Signal transduction: two-component system.	AB0105		σ^7	σ^2
	AB0106		σ^7	σ^2

^aThe functions of the encoded proteins and the AB numbers are indicated according to Miller et al. [15].
doi:10.1371/journal.pone.0044796.t003

ferase are the most down-regulated. Although it does not affect the transcription of its linked TonB-dependent receptor gene (AB2299), σ^7 activated two other TonB-dependent receptor genes AB0705 and AB1870.

The A. butzleri ECF σ Factors Regulate Parts of the Electron Transport Chain

Unlike the *tonB* genes which also appear to be regulated by ECF sigma factors in other bacterial species, we found that a large number of genes regulated by A. butzleri ECF σ factors are involved in the electron transport chain. To verify that the ECF σ factors indeed influence the electron transport chain, we first performed real-time RT-PCR on the ECF-dependent *tonB*, *nap*, *sox* and *tct* genes. We could confirm that the *tonB* genes AB0988, AB1462 and AB1573 are indeed up-regulated by ECF σ^1 , σ^4 and σ^5 respectively (Figure 2a, c and d). ECF σ^2 activates the *nap* and *sox* genes (Figure 2b) of which the latter are repressed by σ^7 (Figure 2e). The real-time RT-PCR results also confirmed that the

tct genes are down-regulated by ECF σ^2 and up-regulated by σ^5 and σ^7 (Figure 2b, d and e).

To prove that the differences in *nap*, *sox* and *tct* transcripts result in phenotypic differences, we determined the nitrate reductase activity, the ability to oxidize sulfate, and the utilization of citrate in ECF mutants compared to the wild-type. As expected, none of the strains except A. butzleri $\Delta\sigma^2::Km$ produced nitrite from the supplied nitrate in the presence of oxygen (Figure 3a). The constitutive expression of σ^2 in the case of $\Delta\sigma^2::Km$ mutant led to nitrite production regardless of the presence of oxygen, indicating that ECF σ^2 up-regulates the Nap-type nitrate reductase. To investigate whether the altered transcription levels of the *sox* genes resulted in a change in Sox enzyme activity, all the strains were grown for 6 hours in AM1 medium supplemented with sulfite and the pH indicator phenol red. The mutants $\Delta\sigma^2$ and $\Delta\sigma^7/\Delta\sigma^7$ further induced the change in medium pH observed during growth of the parental strain, indicating that these mutants have an increased ability to oxidize reduced sulfur compounds




Figure 2. Real-time RT-PCR data confirming parts of the *A. butzleri* ECF regulons. (A) σ^1 , (B) σ^2 , (C) σ^4 , (D) σ^5 , and (E) σ^7 regulon. Transcripts of each $\sigma/A\sigma$ mutant were compared to the cognate $A\sigma$ mutant. Fold change relative to the transcription levels was calculated using the arithmetic formula $2^{-\Delta\Delta Ct}$. Data represent the mean values and standard deviation of four independent experiments with two independent RNA preparations. doi:10.1371/journal.pone.0044796.g002

(Figure 3b). These results confirmed that ECF σ^2 and σ^7 regulate the Sox sulfur oxidation proteins in an opposite way. Finally, we investigated whether these strains are able to utilize citrate, which causes an increase in the pH and a color change of the medium when these bacteria are grown on Simmons agar. In agreement with microarray and real-time RT PCR results, all strains except the mutants $\Delta A\sigma^2$ and $\Delta\sigma^7/A\sigma^7$ caused a color change of the media, indicating that the $\Delta A\sigma^2$ and $\Delta\sigma^7/A\sigma^7$ mutants could not metabolize citrate (Figure 3c).

Discussion

Although *Arcobacter* has been classified as an emerging pathogen (ICMSF, 2002), knowledge on this organism is still limited. The genome sequence of *A. butzleri* strain RM4018 [15] revealed that, unlike other characterized members of the epsilon proteobacteria, this strain possesses seven putative extracytoplasmic function (ECF) σ factors. These σ factors are often involved in the regulation of virulence or stress-related genes in other bacteria. In the present study, we developed genetic tools to manipulate *A. butzleri* and used these tools to investigate the function of the putative RM4018 ECF σ factors. *A. butzleri* is resistant to many different antibiotics [15,29,30]; therefore, it was difficult to find a suitable antibiotic cassette to inactivate the ECF σ factor-encoding genes. Only a kanamycin resistance gene from *C. coli* [31] appeared to be functional in strain RM4018. By electroporation

we were able to generate 10 different mutants and were able to inactivate 5 of the 7 ECF σ factors and to identify the corresponding regulons. Despite repeated attempts, we were unable to mutate the ECF σ factors AB1430 (σ^3) and AB2165 (σ^6), which may indicate that these factors are essential for bacterial growth under the conditions employed.

ECF σ factors are involved in a wide range of environmental responses in many bacterial species [18]. They are mainly divided into stress response ECF σ factors and iron-starvation ECF σ factors [32]. Our results indicate that the ECF σ factors of the strain RM4018 are neither involved in iron-starvation nor in the stress response. Furthermore, we found no evidence of positive regulation of their own transcription, as often found for other ECF σ factors [27]. Except for the *tonB* genes, the RM4018 ECF σ factors regulate completely different sets of genes. This seemingly atypical gene regulation in *Arcobacter* has also been noted for the regulation of the flagellar genes as *Arcobacter* lacks the σ factors FliA and RpoN, which in many bacterial species regulate flagellar biosynthesis [15]. So, in contrast to other eubacteria, many conserved genes in *A. butzleri* appear to be regulated by a σ factor belonging to a different σ factor class; therefore, knowledge on transcription regulation of conserved genes described in other bacterial species cannot be simply extrapolated to *A. butzleri*.

In *A. butzleri* RM4018, 5 out of the 7 ECF σ /anti σ -factor pairs are flanked by genes encoding putative TonB-receptor proteins. In other species, the ECF σ /anti σ factors form, together with an




Figure 3. Functional enzyme assays confirming the differences in transcripts between wild-type and ECF mutants. (A) Nitrate reductase activity was determined under aerobic conditions. Nitrate and nitrite present in the culture supernatant was measured four times and experiments were repeated with two independent cultures. **(B)** Sox enzyme activity assays performed under anaerobic conditions. Standard errors based on four independent experiments are indicated. **(C)** Utilization of citrate tested on Simmons Citrate Agar. Strains which were able to utilize citrate as carbon source caused a change in pH of the Simmons agar which results in a color change of the medium from green to blue. The sign “-” means no bacteria inoculated. doi:10.1371/journal.pone.0044796.g003

outer membrane TonB-dependent receptor, a cell-surface signaling (CSS) system [33]. Transcriptome analyses showed that σ^1 , σ^4 and σ^5 regulate the expression of the TonB-dependent receptors encoded by the same loci, suggesting the putative presence of three potential trans-envelope signal transduction systems in *A. butzleri* RM4018. Most of the described CSS systems are involved in iron signaling and transport [34]. However, other functions have been described including for the *Ralstonia solanacearum* Prha-PrhIR

system, which senses the presence of a plant cell-wall structure and initiates a regulatory cascade that induces the hypersensitive response and the transcription of pathogenicity genes [35]. The role of the three putative CSS systems in *A. butzleri* RM4018 remains to be elucidated, as the growth of mutants in these systems are not affected by iron limitation.

Transcriptome analysis, as well as functional assays, showed that *A. butzleri* ECF σ factors also regulate a number of genes involved

in the binding and transport of specific compounds, energy metabolism and sulfur oxidation. We showed that *Arcobacter* does not reduce nitrate under aerobic conditions (Figure 3A), although it does under anaerobic conditions (data not shown). The ECF σ^2 is involved in this environmental adaptation as the $\Delta\sigma^2::Km$ mutant produces nitrite under aerobic conditions. This may indicate that oxygen stress has an effect on ECF σ^2 activity. Several genes must be indirectly regulated by the ECF σ factors as they were up-regulated in the ECF σ mutants. Interestingly and also seen in *Bacillus subtilis* [36], a number of these genes are regulated by more than one ECF σ factor, which indicates that they may be needed under different growth conditions. For example, σ^2 and σ^7 regulate the *dctPQM* and *sox* genes in an opposite way. The DctPQM proteins form a tripartite ATP-independent periplasmic transporter, which catalyzes the uptake of C₄-dicarboxylates like malate, fumarate and succinate in many aerobic bacteria [37]. The Sox proteins which are not found in other members of the *Campylobacteraceae* are involved in the oxidation of reduced sulfur compounds in sulfur, photo- and chemo-lithotrophic bacteria [38]. Similarly, we showed that σ^5 and σ^7 activate, while σ^2 caused a down-regulation of the expression of the *tctABC* genes encoding a tricarboxylic transport system. Despite this, only σ^7 and σ^2 appeared to have a significant effect on citrate utilization. In many bacterial species a two-component signal transduction system activated by citrate is responsible for the activation of the *tctABC* genes [39]. In *A. butzleri*, the putative TctABC system may also depend on the two-component system (*AB0105–0106*) located directly downstream of the putative *tctABC* genes. The expression of *AB0105–0106* was not dramatically affected by σ^5 and this may explain why citrate utilization was distinct between $\Delta\sigma^5/\Delta\sigma^5$ and $\Delta\sigma^7/\Delta\sigma^7$. All together, the genes regulated by the different *A. butzleri* ECF σ factors indicate that the ECF σ factors form a complex network of regulons that have a major role in regulating bacterial metabolism.

In conclusion, we have shown in this initial study that the ECF σ factors of *A. butzleri* control the transcription of a complex network of regulons that contain genes that are not commonly regulated by ECF σ factors family of proteins. These genes are mainly involved in the energy metabolism. In contrast to other

eubacteria, many conserved genes in *A. butzleri* appear to be regulated by a σ factor belonging to a different σ factor class.

Supporting Information

Table S1 Genes identified by micro-array analyses which are more than fourfold up or down regulated by *A. butzleri* ECF sigma 1.

(DOC)

Table S2 Genes identified by micro-array analyses which are more than fourfold up or down regulated by *A. butzleri* ECF sigma 2.

(DOC)

Table S3 Genes identified by micro-array analyses which are more than fourfold up or down regulated by *A. butzleri* ECF sigma 4.

(DOC)

Table S4 Genes identified by micro-array analyses which are more than fourfold up or down regulated by *A. butzleri* ECF sigma 5.

(DOC)

Table S5 Genes identified by micro-array analyses which are more than fourfold up or down regulated by *A. butzleri* ECF sigma 7.

(DOC)

Acknowledgments

We thank Luis Collado for technical assistance.

Author Contributions

Conceived and designed the experiments: IMM CTP MMSMW. Performed the experiments: IMM RM SH LD IH. Analyzed the data: IMM CTP MMSMW. Contributed reagents/materials/analysis tools: CTP WGM AFA. Wrote the paper: IMM CTP JvP WG WGM MMSMW.

References

- Levican A, Collado L, Aguilar C, Yustes C, Dieguez AL, et al. (2012) *Arcobacter bivahiorum* sp. nov. and *Arcobacter venerupis* sp. nov., new species isolated from shellfish. *Syst Appl Microbiol* 35(3): 133–138.
- Collado L, Figueras MJ (2011) Taxonomy, epidemiology, and clinical relevance of the genus *Arcobacter*. *Clin Microbiol Rev* 24(1): 174–192.
- Neill SD, Ellis WA, O'Brien JJ (1979) Designation of aerotolerant *Campylobacter*-like organisms from porcine and bovine abortions to the genus *Campylobacter*. *Res Vet Sci* 27(2): 180–186.
- On SL, Jensen TK, Bille-Hansen V, Jorsal SE, Vandamme P (2002) Prevalence and diversity of *Arcobacter* spp. isolated from the internal organs of spontaneous porcine abortions in Denmark. *Vet Microbiol* 85(2): 159–167.
- Anderson KF, Kiehlbauch JA, Anderson DC, McClure HM, Wachsmuth IK (1993) *Arcobacter* (*Campylobacter*) *butzleri*-associated diarrheal illness in a nonhuman primate population. *Infect Immun* 61(5): 2220–2223.
- Higgins R, Messier S, Daignault D, Lorange M (1999) *Arcobacter butzleri* isolated from a diarrhoeic non-human primate. *Lab Anim* 33(1): 87–90.
- Lerner J, Brumberger V, Preac-Mursic V (1994) Severe diarrhea associated with *Arcobacter butzleri*. *Eur J Clin Microbiol Infect Dis* 13(8): 660–662.
- Samie A, Obi CL, Barrett LJ, Powell SM, Guerrant RL (2007) Prevalence of *Campylobacter* species, *Helicobacter pylori* and *Arcobacter* species in stool samples from the Venda region, Limpopo, South Africa: Studies using molecular diagnostic methods. *J Infect* 54(6): 558–566.
- Wybo I, Breynaert J, Lauwers S, Lindenburg F, Houf K (2004) Isolation of *Arcobacter skirrovii* from a patient with chronic diarrhea. *J Clin Microbiol* 42(4): 1851–1852.
- Prouzet-Maulon V, Labadi L, Bouges N, Menaud A, Megraud F (2006) *Arcobacter butzleri*: underestimated enteropathogen. *Emerg Infect Dis* 12(2): 307–309.
- ICMSF. (2002) Microorganisms in foods. In: Anonymous Microbiological Testing in Food Safety Management. New York: Kluwer/Plenum: 175.
- Musmanno RA, Russi M, Lior H, Figura N (1997) *In vitro* virulence factors of *Arcobacter butzleri* strains isolated from superficial water samples. *New Microbiol* 20(1): 63–68.
- Rice EW, Rodgers MR, Wesley IV, Johnson CH, Tanner SA (1999) Isolation of *Arcobacter butzleri* from ground water. *Lett Appl Microbiol* 28(1): 31–35.
- Moreno Y, Botella S, Alonso JL, Ferrus MA, Hernandez M, et al. (2003) Specific detection of *Arcobacter* and *Campylobacter* strains in water and sewage by PCR and fluorescent *in situ* hybridization. *Appl Environ Microbiol* 69(2): 1181–1186.
- Miller WG, Parker CT, Rubenfield M, Mendz GL, Wösten MM, et al. (2007) The complete genome sequence and analysis of the epsilonproteobacterium *Arcobacter butzleri*. *PLoS One* 2(12): e1358.
- Helmann JD (2002) The extracytoplasmic function (ECF) sigma factors. *Adv Microb Physiol* 46: 47–110.
- Llamas MA, van der Sar A, Chu BC, Sparrius M, Vogel HJ, et al. (2009) A novel extracytoplasmic function (ECF) sigma factor regulates virulence in *Pseudomonas aeruginosa*. *PLoS Pathog* 5(9): e1000572.
- Shaw LN, Lindholm C, Prajsnar TK, Miller HK, Brown MC, et al. (2008) Identification and characterization of sigma, a novel component of the *Staphylococcus aureus* stress and virulence responses. *PLoS One* 3(12): e3844.
- Raivio TL, Silhavy TJ (2001) Periplasmic stress and ECF sigma factors. *Annu Rev Microbiol* 55: 591–624.
- Campagne S, Damberger FF, Kaczmarczyk A, Francez-Charlot A, Allain FH, et al. (2012) Structural basis for sigma factor mimicry in the general stress response of alphaproteobacteria. *Proc Natl Acad Sci U S A* 109(21): E1405–14.
- Wösten MM (1997) Initiation of transcription and gene organization in *Campylobacter jejuni*. : 120.

22. Yang YH, Speed T (2002) Design issues for cDNA microarray experiments. *Nat Rev Genet* 3(8): 579–588.
23. Lucchini S, Liu H, Jin Q, Hinton JC, Yu J (2005) Transcriptional adaptation of *Shigella flexneri* during infection of macrophages and epithelial cells: Insights into the strategies of a cytosolic bacterial pathogen. *Infect Immun* 73(1): 88–102.
24. Wösten MM, Parker CT, van Mourik A, Guilhabert MR, van Dijk L, et al. (2006) The *Campylobacter jejuni* PhosS/PhosR operon represents a non-classical phosphate-sensitive two-component system. *Mol Microbiol* 62(1): 278–291.
25. Schmittgen TD (2001) Real-time quantitative PCR. *Methods* 25(4): 383–385.
26. Sellars MJ, Hall SJ, Kelly DJ (2002) Growth of *Campylobacter jejuni* supported by respiration of fumarate, nitrate, nitrite, trimethylamine-N-oxide, or dimethyl sulfoxide requires oxygen. *J Bacteriol* 184(15): 4187–4196.
27. Alba BM, Gross CA (2004) Regulation of the *Escherichia coli* sigma-dependent envelope stress response. *Mol Microbiol* 52(3): 613–619.
28. Ho HT, Lipman LJ, Wösten MM, van Asten AJ, Gastra W (2008) *Arcobacter* spp. possess two very short flagellins of which FlaA is essential for motility. *FEMS Immunol Med Microbiol* 53(1): 85–95.
29. Atabay HI, Aydin F (2001) Susceptibility of *Arcobacter butzleri* isolates to 23 antimicrobial agents. *Lett Appl Microbiol* 33(6): 430–433.
30. Houf K, Devriese LA, Haesebrouck F, Vandenberg O, Butzler JP, et al. (2004) Antimicrobial susceptibility patterns of *Arcobacter butzleri* and *Arcobacter cryaerophilus* strains isolated from humans and broilers. *Microb Drug Resist* 10(3): 243–247.
31. Labigne-Roussel A, Harel J, Tompkins L (1987) Gene transfer from *Escherichia coli* to *Campylobacter* species: Development of shuttle vectors for genetic analysis of *Campylobacter jejuni*. *J Bacteriol* 169(11): 5320–3.
32. Braun V, Mahren S, Ogierman M (2003) Regulation of the FecI-type ECF sigma factor by transmembrane signalling. *Curr Opin Microbiol* 6(2): 173–180.
33. Visca P, Leoni L, Wilson MJ, Lamont IL (2002) Iron transport and regulation, cell signalling and genomics: Lessons from *Escherichia coli* and *Pseudomonas*. *Mol Microbiol* 45(5): 1177–1190.
34. Koebnik R (2005) TonB-dependent trans-envelope signalling: The exception or the rule? *Trends Microbiol* 13(8): 343–347.
35. Brito B, Aldon D, Barberis P, Boucher C, Genin S (2002) A signal transfer system through three compartments transduces the plant cell contact-dependent signal controlling *Ralstonia solanacearum* hrp genes. *Mol Plant Microbe Interact* 15(2): 109–119.
36. Luo Y, Asai K, Sadaie Y, Helmann JD (2010) Transcriptomic and phenotypic characterization of a *Bacillus subtilis* strain without extracytoplasmic function σ factors. *J Bacteriol* 192: 5736–45.
37. Janausch IG, Zientz E, Tran QH, Kroger A, Uden G (2002) C4-dicarboxylate carriers and sensors in bacteria. *Biochim Biophys Acta* 1553(1–2): 39–56.
38. Ghosh W, Dam B (2009) Biochemistry and molecular biology of lithotrophic sulfur oxidation by taxonomically and ecologically diverse bacteria and archaea. *FEMS Microbiol Rev* 33(6): 999–1043.
39. Brocker M, Schaffer S, Mack C, Bott M (2009) Citrate utilization by *Corynebacterium glutamicum* is controlled by the CitAB two-component system through positive regulation of the citrate transport genes *citH* and *tetCBA*. *J Bacteriol* 191(12): 3869–3880.